

Evolution, Physiology and Phytochemistry of the Psychotoxic Arable Mimic Weed Darnel (*Lolium temulentum* L.)

Howard Thomas, Jayne Elisabeth Archer, and Richard Marggraf Turley

Contents

1	Introduction	74
2	Phylogeny and Evolution of <i>L. temulentum</i>	74
2.1	The <i>L. temulentum</i> Genome	74
2.2	Molecular Systematics of <i>Lolium</i>	76
2.3	Selection of Domestication Traits in <i>L. temulentum</i>	77
2.4	The Spread and Perpetuation of <i>L. temulentum</i> in Cereal Grain Stocks	79
2.5	Other <i>Lolium</i> spp. Following a Similar Evolutionary Pathway	79
3	Physiology and Biochemistry of <i>L. temulentum</i>	81
3.1	A Model for the Study of Photoperiodic Control of Flowering	81
3.2	Leaf Development and Senescence in <i>L. temulentum</i>	83
3.3	Carbohydrate Metabolism and Carbon Partitioning	85
3.4	<i>L. temulentum</i> and Abiotic Stress	88
4	Biotic Interactions and Toxicity of <i>L. temulentum</i>	91
4.1	Symptoms of Darnel Poisoning	91
4.2	Fungal Endophytes and the Chemistry of <i>Lolium</i> Toxins	92
4.3	Darnel and Ergot	93
4.4	Other Possible Sources of Toxicity in <i>L. temulentum</i>	94
5	<i>L. temulentum</i> in History and Literature	95
	References	97

Abstract Darnel (*Lolium temulentum* L.), the subject of this review, is botanically and culturally significant because of its evolutionary origin as a mimic weed of cereals and its reputation as a source of potent psychoactive toxins. Evidence from molecular phylogeny, palaeontology and archaeology allows the source and spread

H. Thomas (✉)

IBERS, Edward Llwyd Building, Penglais Campus, Aberystwyth University, Ceredigion SY23 3DA, UK
hot@aber.ac.uk

J.E. Archer and R. Marggraf Turley

Department of English and Creative Writing, Penglais Campus, Aberystwyth University,
Ceredigion SY23 3DY, UK